

VARIATIONS

In this unit, students will learn how to

- ✎ *define ratio, proportions and variations (direct and inverse).*
- ✎ *find 3rd, 4th, mean and continued proportion.*
- ✎ *apply theorems of invertendo, alternendo, componendo, dividendo and componendo & dividendo to find proportions.*
- ✎ *define joint variation.*
- ✎ *solve problems related to joint variation.*
- ✎ *use k-method to prove conditional equalities involving proportions.*
- ✎ *solve real life problems based on variations.*

3.1 Ratio, Proportions and Variations

3.1(i) Define (a) ratio, (b) proportion and (c) variations (direct and inverse).

(a) Ratio

A relation between two quantities of the same kind (measured in same unit) is called **ratio**. If a and b are two quantities of the same kind and b is not zero, then the ratio of a and b is written as $a : b$ or in fraction $\frac{a}{b}$

e.g., if a hockey team wins 4 games and loses 5, then the ratio of the games won to games lost is $4 : 5$ or in fraction $\frac{4}{5}$

Remember that:

- (i) The **order** of the elements in a ratio is important.
- (ii) In ratio $a : b$, the first term a is called **antecedent** and the second term b is called **consequent**.
- (iii) A ratio has no **units**.

Example 1: Find the ratio of

- (i) 200gm to 700 gm
- (ii) 1km to 600m

Solution: (i) Ratio of 200gm to 700 gm

$$200 : 700 = \frac{200}{700} = \frac{2}{7} = 2 : 7$$

Where $2 : 7$ is the simplest (lowest) form of the ratio $200 : 700$.

- (ii) Ratio of 1km to 600m

Since $1\text{km} = 1000\text{m}$

then $1000 : 600 = \frac{1000}{600} = \frac{10}{6} = \frac{5}{3} = 5 : 3$

or $1\text{km} : 600\text{m} = 1000:600$

$$= \frac{1000}{100} : \frac{600}{100} = 10:6 = 5:3$$

Example 2: Find a , if the ratios $a + 3 : 7 + a$ and $4 : 5$ are equal.

Solution: Since the ratios $a + 3 : 7 + a$ and $4 : 5$ are equal.

\therefore in fraction form

$$\frac{a + 3}{7 + a} = \frac{4}{5}$$

$$5(a + 3) = 4(7 + a)$$

$$5a + 15 = 28 + 4a$$

$$5a - 4a = 28 - 15$$

$$a = 13$$

Thus the given ratios will be equal if $a = 13$.

Example 3: If 2 is added in each number of the ratio 3 : 4, we get a new ratio 5 : 6. Find the numbers.

Solution: Because the ratio of two numbers is 3 : 4.

Multiply each number of the ratio with x . Then the numbers be $3x$, $4x$ and the ratio becomes $3x:4x$. Now according to the given condition

$$\frac{3x + 2}{4x + 2} = \frac{5}{6}$$

$$6(3x + 2) = 5(4x + 2) \Rightarrow 18x + 12 = 20x + 10$$

$$18x - 20x = 10 - 12 \Rightarrow -2x = -2 \Rightarrow x = 1$$

Thus the required numbers are

$$3x = 3(1) = 3$$

and $4x = 4(1) = 4$.

Example 4: Find the ratio $3a + 4b : 5a + 7b$ if $a : b = 5 : 8$.

Solution: Given that $a : b = 5 : 8$ or $\frac{a}{b} = \frac{5}{8}$

Now $3a + 4b : 5a + 7b = \frac{3a + 4b}{5a + 7b}$

$$= \frac{\frac{3a + 4b}{b}}{\frac{5a + 7b}{b}} = \frac{3\left(\frac{a}{b}\right) + 4\left(\frac{b}{b}\right)}{5\left(\frac{a}{b}\right) + 7\left(\frac{b}{b}\right)} \quad (\text{Dividing numerator and denominator by } b)$$

$$= \frac{3\left(\frac{5}{8}\right) + 4(1)}{5\left(\frac{5}{8}\right) + 7(1)} \quad \left(\because \frac{a}{b} = \frac{5}{8}\right)$$

$$= \frac{\frac{15}{8} + 4}{\frac{25}{8} + 7} = \frac{\frac{15 + 32}{8}}{\frac{25 + 56}{8}} = \frac{47}{81}$$

Hence, $3a + 4b : 5a + 7b = 47 : 81$.

(b) Proportion

A **proportion** is a statement, which is expressed as an equivalence of two ratios.

If two ratios $a : b$ and $c : d$ are equal, then we can write $a : b = c : d$

Where quantities a, d are called **extremes**, while b, c are called **means**.

Symbolically the proportion of a, b, c and d is written as

$$a : b :: c : d$$

or $a : b = c : d$

$$\text{or } \frac{a}{b} = \frac{c}{d}$$

$$\text{i.e., } ad = bc$$

This shows that, **Product of extremes = Product of means.**

Example 5: Find x , if $60\text{m} : 90\text{m} :: 20\text{kg} : x \text{ kg}$

Solution: Given that $60\text{m} : 90\text{m} :: 20\text{kg} : x \text{ kg}$
 $60 : 90 = 20 : x$

\therefore Product of extremes = Product of means

$$\therefore 60x = 90 \times 20$$

$$x = \frac{90 \times 20}{60} = 30 \text{ i.e., } x \text{ is } 30 \text{ kg}$$

Example 6: Find the cost of 15kg of sugar, if 7 kg of sugar costs 560 rupees.

Solution: Let the cost of 15kg of sugar be x -rupees.

Then in proportion form

$$15\text{kg} : 7\text{kg} :: \text{Rs. } x : \text{Rs. } 560$$

$$15 : 7 = x : 560$$

\therefore Product of extremes = Product of means

$$\therefore 15 \times 560 = 7x$$

$$7x = 15 \times 560$$

$$x = \frac{15 \times 560}{7} = 15(80) = 1200$$

Thus, $x = \text{Rs. } 1200$.

EXERCISE 3.1

- Express the following as a ratio $a : b$ and as a fraction in its simplest (lowest) form.
 - Rs. 750 , Rs. 1250
 - 450cm , 3 m
 - 4kg , 2kg 750gm
 - 27min. 30 sec, 1 hour
 - 75° , 225°
- In a class of 60 students, 25 students are girls and remaining students are boys. Compute the ratio of
 - boys to total students
 - boys to girls
- If $3(4x - 5y) = 2x - 7y$, find the ratio $x : y$.
- Find the value of p , if the ratios $2p + 5 : 3p + 4$ and $3 : 4$ are equal.
- If the ratios $3x + 1 : 6 + 4x$ and $2 : 5$ are equal. Find the value of x .

6. Two numbers are in the ratio 5 : 8. If 9 is added to each number, we get a new ratio 8 : 11. Find the numbers.
7. If 10 is added in each number of the ratio 4 : 13, we get a new ratio 1 : 2. What are the numbers?
8. Find the cost of 8kg of mangoes, if 5kg of mangoes cost Rs. 250.
9. If $a : b = 7 : 6$, find the value of $3a + 5b : 7b - 5a$.
10. Complete the following:
- (i) If $\frac{24}{7} = \frac{6}{x}$, then $4x =$ _____
- (ii) If $\frac{5a}{3x} = \frac{15b}{y}$, then $ay =$ _____
- (iii) If $\frac{9pq}{2lm} = \frac{18p}{5m}$, then $5q =$ _____
11. Find x in the following proportions.
- (i) $3x - 2 : 4 :: 2x + 3 : 7$ (ii) $\frac{3x - 1}{7} : \frac{3}{5} :: \frac{2x}{3} : \frac{7}{5}$
- (iii) $\frac{x - 3}{2} : \frac{5}{x - 1} :: \frac{x - 1}{3} : \frac{4}{x + 4}$ (iv) $p^2 + pq + q^2 : x :: \frac{p^3 - q^3}{p + q} : (p - q)^2$
- (v) $8 - x : 11 - x :: 16 - x : 25 - x$

(c) Variation:

The word variation is frequently used in all sciences. There are two types of variations: (i) Direct variation (ii) Inverse variation.

(i) Direct Variation

If two quantities are related in such a way that increase (decrease) in one quantity causes increase (decrease) in the other quantity, then this variation is called **direct variation**.

In other words, if a quantity y varies directly with regard to a quantity x . We say that y is **directly proportional** to x and is written as $y \propto x$ or $y = kx$. i.e., $\frac{y}{x} = k, k \neq 0$.

The sign \propto read as “varies as” is called the sign of proportionality or variation, while $k \neq 0$ is known as constant of variation.

e.g., (i) Faster the speed of a car, longer the distance it covers.

(ii) The smaller the radius of the circle, smaller the circumference is.

Example 1: Find the relation between distance d of a body falling from rest varies directly as the square of the time t , neglecting air resistance. Find k , if $d = 16$ feet for $t = 1$ sec. Also derive a relation between d and t .

Solution: Since d is the distance of the body falling from rest in time t .

Then under the given condition

$$d \propto t^2$$

$$\text{i.e., } d = kt^2 \quad \text{(i)}$$

Since $d = 16$ feet and $t = 1$ sec

Then equation (i) becomes

$$16 = k(1)^2$$

$$\text{i.e., } k = 16$$

put in eq. (i) $d = 16t^2$

Which is a relationship between the distance d and time t .

Activity:

From the above example:

- (i) Find time t , when $d = 64$ feet
- (ii) Find distance d , when $t = 3$ sec

Example 2: If y varies directly as x , find

- (a) the equation connecting x and y .
- (b) the constant of variation k and the relation between x and y , when $x = 7$ and $y = 6$
- (c) the value of y , when $x = 21$.

Solution: (a) Given that y varies directly as x .

Therefore $y \propto x$, i.e., $y = kx$, where k is constant of variation.

(b) Putting $x = 7$ and $y = 6$ in equation

$$y = kx \quad \text{(i)}$$

$$\text{We get } 6 = 7k \Rightarrow k = \frac{6}{7}$$

$$\text{Put in eq. (i) } y = \frac{6}{7}x \quad \text{(ii)}$$

(c) Now put $x = 21$, in equation (ii)

$$\text{Then } y = \frac{6}{7}(21) = 18$$

Example 3: Given that A varies directly as the square of r and $A = \frac{1782}{7}$ cm², when $r = 9$ cm.

If $r = 14$ cm, then find A .

Solution: Since A varies directly as square of r

$$\therefore A \propto r^2$$

$$\text{or } A = kr^2 \quad \text{(i)}$$

$$\frac{1782}{7} = k(9)^2$$

$$\frac{1782}{7 \times 81} = k \quad \text{or} \quad k = \frac{22}{7}$$

Put $k = \frac{22}{7}$ and $r = 14\text{cm}$ in eq. (i)

$$A = \frac{22}{7} (14)^2 = \frac{22}{7} \times 14 \times 14 = 616 \text{ cm}^2$$

Example 4: If y varies directly as cube of x and $y = 81$ when $x = 3$, so evaluate y when $x = 5$.

Solution: Given that y varies directly as cube of x .

$$\text{i.e., } y \propto x^3 \quad \text{or} \quad y = kx^3 \quad (\text{i}) \quad (\text{where } k \text{ is constant})$$

Put $y = 81$ and $x = 3$ in (i)

$$81 = k(3)^3 \Rightarrow 27k = 81 \Rightarrow k = 3$$

Now put $k = 3$ and $x = 5$ in eq. (i)

$$y = 3(5)^3 = 375$$

(ii) Inverse Variation

If two quantities are related in such a way that when one quantity increases, the other decreases is called **inverse variation**.

In other words, if a quantity y varies inversely with regard to quantity x . We say that y is inversely proportional to x or y varies inversely as x and is written as $y \propto \frac{1}{x}$ or $y = \frac{k}{x}$.

i.e., $xy = k$, where $k \neq 0$ is the constant of variation.

Example 1: If y varies inversely as x and $y = 8$, when $x = 4$. Find y , when $x = 16$.

Solution: Since y varies inversely as x , therefore

$$y \propto \frac{1}{x} \quad \text{or} \quad y = \frac{k}{x} \quad (\text{i})$$

$$\Rightarrow xy = k \quad (\text{ii})$$

Putting $y = 8$ and $x = 4$ in (ii)

$$\begin{aligned} k &= (x)(y) \\ &= (4)(8) = 32 \end{aligned}$$

Now put $k = 32$ and $x = 16$ in (i) $\Rightarrow y = \frac{32}{16} = 2$

Example 2: If y varies inversely as x^2 and $y = 16$, when $x = 5$, so find x , when $y = 100$.

Solution: Since y varies inversely as x^2 , therefore

$$\begin{aligned} y &\propto \frac{1}{x^2} \quad \text{or} \quad y = \frac{k}{x^2} \\ k &= x^2y \quad (\text{i}) \end{aligned}$$

Put $x = 5$ and $y = 16$ in (i)

$$k = (5)^2 \times 16$$

$$k = 400$$

Now put $k = 400$ and $y = 100$ in (i)

$$400 = 100x^2 \quad \text{or} \quad x^2 = \frac{400}{100} = 4$$

$$x = \pm 2$$

EXERCISE 3.2

1. If y varies directly as x , and $y = 8$ when $x = 2$, find
 - (i) y in terms of x
 - (ii) y when $x = 5$
 - (iii) x when $y = 28$
2. If $y \propto x$, and $y = 7$ when $x = 3$ find
 - (i) y in terms of x
 - (ii) x when $y = 35$ and y when $x = 18$
3. If $R \propto T$ and $R = 5$ when $T = 8$, find the equation connecting R and T . Also find R when $T = 64$ and T when $R = 20$.
4. If $R \propto T^2$ and $R = 8$ when $T = 3$, find R when $T = 6$.
5. If $V \propto R^3$ and $V = 5$ when $R = 3$, find R when $V = 625$.
6. If w varies directly as u^3 and $w = 81$ when $u = 3$. Find w when $u = 5$.
7. If y varies inversely as x and $y = 7$ when $x = 2$, find y when $x = 126$.
8. If $y \propto \frac{1}{x}$ and $y = 4$ when $x = 3$, find x when $y = 24$.
9. If $w \propto \frac{1}{z}$ and $w = 5$ when $z = 7$, find w when $z = \frac{175}{4}$.
10. $A \propto \frac{1}{r^2}$ and $A = 2$ when $r = 3$, find r when $A = 72$.
11. $a \propto \frac{1}{b^2}$ and $a = 3$ when $b = 4$, find a when $b = 8$.
12. $V \propto \frac{1}{r^3}$ and $V = 5$ when $r = 3$, find V when $r = 6$ and r when $V = 320$.
13. $m \propto \frac{1}{n^3}$ and $m = 2$ when $n = 4$, find m when $n = 6$ and n when $m = 432$.

3.1(ii) Find 3rd, 4th, mean and continued proportion:

We are already familiar with proportions that if quantities a , b , c and d are in proportion, then $a : b :: c : d$

i.e., product of extremes = product of means

Third Proportional

If three quantities a , b and c are related as $a : b :: b : c$, then c is called the third proportion.

Example 1: Find a third proportional of $x + y$ and $x^2 - y^2$.

Solution: Let c be the third proportional,

$$\text{then } x + y : x^2 - y^2 :: x^2 - y^2 : c$$

$$c(x + y) = (x^2 - y^2)(x^2 - y^2)$$

$$c = \frac{(x^2 - y^2)(x^2 - y^2)}{x + y} = \frac{(x^2 - y^2)(x - y)(x + y)}{(x + y)}$$

$$c = (x^2 - y^2)(x - y) = (x + y)(x - y)^2$$

Fourth Proportional

If four quantities a , b , c and d are related as

$$a : b :: c : d$$

Then d is called the fourth proportional.

Example 2: Find fourth proportional of $a^3 - b^3$, $a + b$ and $a^2 + ab + b^2$

Solution: Let x be the fourth proportional,

$$\text{then } (a^3 - b^3) : (a + b) :: (a^2 + ab + b^2) : x$$

$$\text{i.e., } x(a^3 - b^3) = (a + b)(a^2 + ab + b^2)$$

$$x = \frac{(a + b)(a^2 + ab + b^2)}{a^3 - b^3} = \frac{(a + b)(a^2 + ab + b^2)}{(a - b)(a^2 + ab + b^2)}$$

$$x = \frac{a + b}{a - b}$$

Mean Proportional

If three quantities a , b and c are related as $a : b :: b : c$,

then b is called the mean proportional.

Example 3: Find the mean proportional of $9p^6q^4$ and r^8 .

Solution: Let m be the mean proportional,

$$\text{then } 9p^6q^4 : m :: m : r^8$$

$$\text{or } m \cdot m = 9p^6q^4(r^8)$$

$$m^2 = 9p^6q^4r^8$$

$$m = \pm \sqrt{9p^6q^4r^8} = \pm 3p^3q^2r^4$$

Continued Proportion

If three quantities a , b and c are related as

$$a : b :: b : c,$$

where a is first, b is the mean and c is the third proportional, then a , b and c are in continued proportion.

Example 4: Find p , if 12, p and 3 are in continued proportion.

Solution: Since 12, p and 3 are in continued proportion.

$$\therefore 12 : p :: p : 3 \quad \text{i.e., } p \cdot p = (12)(3) \Rightarrow p^2 = 36$$

$$\text{Thus, } p = \pm 6$$

EXERCISE 3.3

1. Find a third proportional to

(i) 6, 12	(ii) $a^3, 3a^2$
(iii) $a^2 - b^2, a - b$	(iv) $(x - y)^2, x^3 - y^3$
(v) $(x + y)^2, x^2 - xy - 2y^2$	(vi) $\frac{p^2 - q^2}{p^3 + q^3}, \frac{p - q}{p^2 - pq + q^2}$

2. Find a fourth proportional to

(i) 5, 8, 15	(ii) $4x^4, 2x^3, 18x^5$
(iii) $15a^5b^6, 10a^2b^5, 21a^3b^3$	(iv) $x^2 - 11x + 24, (x - 3), 5x^4 - 40x^3$
(v) $p^3 + q^3, p^2 - q^2, p^2 - pq + q^2$	
(vi) $(p^2 - q^2)(p^2 + pq + q^2), p^3 + q^3, p^3 - q^3$	

3. Find a mean proportional between

(i) 20, 45	(ii) $20x^3y^5, 5x^7y$
(iii) $15p^4qr^3, 135q^5r^7$	(iv) $x^2 - y^2, \frac{x - y}{x + y}$

4. Find the values of the letter involved in the following continued proportions.

(i) 5, p , 45	(ii) 8, x , 18
(iii) 12, $3p - 6$, 27	(iv) 7, $m - 3$, 28

3.2 Theorems on Proportions

If four quantities a, b, c and d form a proportion, then many other useful properties may be deduced by the properties of fractions.

(1) Theorem of Invertendo

If $a : b = c : d$, then $b : a = d : c$

Example 1: If $3m : 2n = p : 2q$, then

$$2n : 3m = 2q : p$$

Solution: Since $3m : 2n = p : 2q$

$$\therefore \frac{3m}{2n} = \frac{p}{2q}$$

By invertendo theorem

$$\frac{2n}{3m} = \frac{2q}{p}$$

$$\text{i.e., } 2n : 3m = 2q : p$$

(2) Theorem of Alternando

If $a : b = c : d$, then $a : c = b : d$

Example 2: If $3p + 1 : 2q = 5r : 7s$, then prove that $3p + 1 : 5r = 2q : 7s$

Solution: Given that $3p + 1 : 2q = 5r : 7s$

Then $\frac{3p+1}{2q} = \frac{5r}{7s}$

By alternando theorem

$$\frac{3p+1}{5r} = \frac{2q}{7s}$$

Thus, $3p+1 : 5r = 2q : 7s$

(3) Theorem of Componendo

If $a : b = c : d$, then

(i) $a + b : b = c + d : d$

and (ii) $a : a + b = c : c + d$

Example 3: If $m + 3 : n = p : q - 2$, then

$$m + n + 3 : n = p + q - 2 : q - 2$$

Solution: Since $m + 3 : n = p : q - 2$

$$\therefore \frac{m+3}{n} = \frac{p}{q-2}$$

By componendo theorem

$$\frac{(m+3)+n}{n} = \frac{p+(q-2)}{q-2}$$

or $\frac{m+n+3}{n} = \frac{p+q-2}{q-2}$

Thus $m + n + 3 : n = p + q - 2 : q - 2$

(4) Theorem of Dividendo

If $a : b = c : d$, then

(i) $a - b : b = c - d : d$

and (ii) $a : a - b = c : c - d$

Example 4: If $m + 1 : n - 2 = 2p + 3 : 3q + 1$.

Then $m - n + 3 : n - 2 = 2p - 3q + 2 : 3q + 1$

Solution: Given that $m + 1 : n - 2 = 2p + 3 : 3q + 1$

Then $\frac{m+1}{n-2} = \frac{2p+3}{3q+1}$

By dividendo theorem

$$\frac{m-n+3}{n-2} = \frac{2p-3q+2}{3q+1}$$

Thus $m - n + 3 : n - 2 = 2p - 3q + 2 : 3q + 1$

(5) Theorem of Componendo-dividendo

If $a : b = c : d$, then

(i) $a + b : a - b = c + d : c - d$

and (ii) $a - b : a + b = c - d : c + d$

Example 5: If $m : n = p : q$.

Then prove that $3m + 7n : 3m - 7n = 3p + 7q : 3p - 7q$

Solution: Since $m : n = p : q$

$$\text{or } \frac{m}{n} = \frac{p}{q}$$

Multiplying both sides by $\frac{3}{7}$, we get

$$\frac{3m}{7n} = \frac{3p}{7q}$$

Then using componendo-dividendo theorem

$$\frac{3m + 7n}{3m - 7n} = \frac{3p + 7q}{3p - 7q}$$

Thus $3m + 7n : 3m - 7n = 3p + 7q : 3p - 7q$

Example 6: If $5m + 3n : 5m - 3n = 5p + 3q : 5p - 3q$,

then show that $m : n = p : q$

Solution: Given that $5m + 3n : 5m - 3n = 5p + 3q : 5p - 3q$

$$\text{or } \frac{5m + 3n}{5m - 3n} = \frac{5p + 3q}{5p - 3q}$$

By componendo-dividendo theorem

$$\frac{(5m + 3n) + (5m - 3n)}{(5m + 3n) - (5m - 3n)} = \frac{(5p + 3q) + (5p - 3q)}{(5p + 3q) - (5p - 3q)}$$

$$\frac{5m + 3n + 5m - 3n}{5m + 3n - 5m + 3n} = \frac{5p + 3q + 5p - 3q}{5p + 3q - 5p + 3q}$$

$$\frac{10m}{6n} = \frac{10p}{6q}$$

Multiplying both sides by $\frac{6}{10}$.

$$\frac{m}{n} = \frac{p}{q}$$

i.e., $m : n = p : q$

Example 7: Using theorem of componendo-dividendo, find the value of

$$\frac{m + 3p}{m - 3p} + \frac{m + 2q}{m - 2q}, \text{ if } m = \frac{6pq}{p + q}.$$

Solution: Since $m = \frac{6pq}{p+q}$ or $m = \frac{(3p)(2q)}{p+q}$ (i)

$$\therefore \frac{m}{3p} = \frac{2q}{p+q}$$

By componendo-dividendo theorem

$$\frac{m+3p}{m-3p} = \frac{2q+(p+q)}{2q-(p+q)} = \frac{2q+p+q}{2q-p-q}$$

$$\frac{m+3p}{m-3p} = \frac{p+3q}{q-p} \quad \text{(ii)}$$

Again from eq. (i), we have

$$\frac{m}{2q} = \frac{3p}{p+q}$$

By componendo-dividendo theorem

$$\frac{m+2q}{m-2q} = \frac{3p+(p+q)}{3p-(p+q)} = \frac{3p+p+q}{3p-p-q}$$

$$\frac{m+2q}{m-2q} = \frac{4p+q}{2p-q} \quad \text{(iii)}$$

Adding (ii) and (iii)

$$\begin{aligned} \frac{m+3p}{m-3p} + \frac{m+2q}{m-2q} &= \frac{p+3q}{q-p} + \frac{4p+q}{2p-q} = -\frac{p+3q}{p-q} + \frac{4p+q}{2p-q} \\ &= \frac{-(p+3q)(2p-q) + (p-q)(4p+q)}{(p-q)(2p-q)} \\ &= \frac{-2p^2 - 5pq + 3q^2 + 4p^2 - 3pq - q^2}{(p-q)(2p-q)} \\ &= \frac{2p^2 - 8pq + 2q^2}{(p-q)(2p-q)} = \frac{2(p^2 - 4pq + q^2)}{(p-q)(2p-q)} \end{aligned}$$

Example 8: Using theorem of componendo-dividendo, solve the equation

$$\frac{\sqrt{x+3} + \sqrt{x-3}}{\sqrt{x+3} - \sqrt{x-3}} = \frac{4}{3}$$

Solution: Given equation is $\frac{\sqrt{x+3} + \sqrt{x-3}}{\sqrt{x+3} - \sqrt{x-3}} = \frac{4}{3}$

By componendo-dividendo theorem

$$\frac{\sqrt{x+3} + \sqrt{x-3} + \sqrt{x+3} - \sqrt{x-3}}{\sqrt{x+3} + \sqrt{x-3} - \sqrt{x+3} + \sqrt{x-3}} = \frac{4+3}{4-3}$$

$$\frac{2\sqrt{x+3}}{2\sqrt{x-3}} = \frac{7}{1} \Rightarrow \sqrt{\frac{x+3}{x-3}} = 7$$

Squaring both sides

$$\frac{x+3}{x-3} = 49$$

$$x+3 = 49(x-3) \Rightarrow x+3 = 49x-147 \Rightarrow x-49x = -147-3$$

$$-48x = -150 \Rightarrow 48x = 150 \Rightarrow x = \frac{150}{48} = \frac{25}{8}$$

Example 9: Using componendo-dividendo theorem, solve the equation $\frac{(x+3)^2 - (x-5)^2}{(x+3)^2 + (x-5)^2} = \frac{4}{5}$.

Solution: Given equation is $\frac{(x+3)^2 - (x-5)^2}{(x+3)^2 + (x-5)^2} = \frac{4}{5}$

By componendo-dividendo theorem

$$\frac{(x+3)^2 - (x-5)^2 + (x+3)^2 + (x-5)^2}{(x+3)^2 - (x-5)^2 - (x+3)^2 - (x-5)^2} = \frac{4+5}{4-5}$$

$$\frac{2(x+3)^2}{-2(x-5)^2} = \frac{9}{-1} \Rightarrow \left(\frac{x+3}{x-5}\right)^2 = (3)^2$$

Taking square root $\frac{x+3}{x-5} = \pm 3$

$$\frac{x+3}{x-5} = 3$$

$$\text{or } \frac{x+3}{x-5} = -3$$

$$x+3 = 3(x-5)$$

$$x+3 = -3(x-5)$$

$$x+3 = 3x-15$$

$$x+3 = -3x+15$$

$$-2x = -18$$

$$4x = 12$$

$$x = 9$$

$$x = 3$$

\therefore The solution set is $\{3, 9\}$

EXERCISE 3.4

1. Prove that $a : b = c : d$, if

(i) $\frac{4a+5b}{4a-5b} = \frac{4c+5d}{4c-5d}$

(ii) $\frac{2a+9b}{2a-9b} = \frac{2c+9d}{2c-9d}$

(iii) $\frac{ac^2+bd^2}{ac^2-bd^2} = \frac{c^3+d^3}{c^3-d^3}$

(iv) $\frac{a^2c+b^2d}{a^2c-b^2d} = \frac{ac^2+bd^2}{ac^2-bd^2}$

(v) $pa+qb : pa-qb = pc+qd : pc-qd$

(vi) $\frac{a+b+c+d}{a+b-c-d} = \frac{a-b+c-d}{a-b-c+d}$

$$(vii) \frac{2a + 3b + 2c + 3d}{2a + 3b - 2c - 3d} = \frac{2a - 3b + 2c - 3d}{2a - 3b - 2c + 3d}$$

$$(viii) \frac{a^2 + b^2}{a^2 - b^2} = \frac{ac + bd}{ac - bd}$$

2. Using theorem of componendo-dividendo

$$(i) \text{ Find the value of } \frac{x + 2y}{x - 2y} + \frac{x + 2z}{x - 2z}, \text{ if } x = \frac{4yz}{y + z}$$

$$(ii) \text{ Find the value of } \frac{m + 5n}{m - 5n} + \frac{m + 5p}{m - 5p}, \text{ if } m = \frac{10np}{n + p}$$

$$(iii) \text{ Find the value of } \frac{x - 6a}{x + 6a} - \frac{x + 6b}{x - 6b}, \text{ if } x = \frac{12ab}{a - b}$$

$$(iv) \text{ Find the value of } \frac{x - 3y}{x + 3y} - \frac{x + 3z}{x - 3z}, \text{ if } x = \frac{3yz}{y - z}$$

$$(v) \text{ Find the value of } \frac{s - 3p}{s + 3p} + \frac{s + 3q}{s - 3q}, \text{ if } s = \frac{6pq}{p - q}$$

$$(vi) \text{ Solve } \frac{(x - 2)^2 - (x - 4)^2}{(x - 2)^2 + (x - 4)^2} = \frac{12}{13}$$

$$(vii) \text{ Solve } \frac{\sqrt{x^2 + 2} + \sqrt{x^2 - 2}}{\sqrt{x^2 + 2} - \sqrt{x^2 - 2}} = 2$$

$$(viii) \text{ Solve } \frac{\sqrt{x^2 + 8p^2} - \sqrt{x^2 - p^2}}{\sqrt{x^2 + 8p^2} + \sqrt{x^2 - p^2}} = \frac{1}{3}$$

$$(ix) \text{ Solve } \frac{(x + 5)^3 - (x - 3)^3}{(x + 5)^3 + (x - 3)^3} = \frac{13}{14}$$

3.3.(i) Joint variation

A combination of direct and inverse variations of one or more than one variables forms **joint variation**.

If a variable y varies directly as x and varies inversely as z .

$$\text{Then } y \propto x \quad \text{and} \quad y \propto \frac{1}{z}$$

In joint variation, we write it as

$$y \propto \frac{x}{z}$$

$$\text{i.e., } y = k \frac{x}{z}$$

Where $k \neq 0$ is the constant of variation.

For example, by Newton's law of gravitation, if one body attracts another with a force (G), that varies directly as the product of their masses (m_1), (m_2) and inversely as the square of the distance (d) between them.

$$\text{i.e., } G \propto \frac{m_1 m_2}{d^2} \text{ or } G = k \frac{m_1 m_2}{d^2}, \text{ where } k \neq 0 \text{ is the constant}$$

3.3.(ii) Problems related to joint variation.

Procedure to solve the problems related to joint variation is explained through examples.

Example 1: If y varies jointly as x^2 and z and $y = 6$ when $x = 4$, $z = 9$. Write y as a function of x and z and determine the value of y , when $x = -8$ and $z = 12$.

Solution: Since y varies jointly as x^2 and z , therefore

$$\text{i.e., } y \propto x^2 z$$

$$y = kx^2 z \quad (\text{i})$$

$$\text{Put } y = 6, x = 4, z = 9$$

$$6 = k(4)^2(9)$$

$$\frac{6}{16 \times 9} = k \Rightarrow k = \frac{1}{24}$$

$$\text{Put } k = \frac{1}{24} \text{ in eq.(i), } y = \frac{1}{24} x^2 z$$

Now put $x = -8$, $z = 12$ in the above equation,

$$y = \frac{1}{24} (-8)^2 (12) = 32$$

Example 2: p varies jointly as q and r^2 and inversely as s and t^2 , $p = 40$, when $q = 8$, $r = 5$, $s = 3$, $t = 2$. Find p in terms of q , r , s and t . Also find the value of p when $q = -2$, $r = 4$, $s = 3$ and $t = -1$.

Solution: Given that $p \propto \frac{qr^2}{st^2}$

$$p = k \frac{qr^2}{st^2} \quad (\text{i})$$

$$\text{Put } p = 40, q = 8, r = 5, s = 3 \text{ and } t = 2$$

$$40 = k \frac{(8)(5)^2}{3(2)^2}$$

$$\frac{40 \times 3 \times 4}{8 \times 25} = k$$

$$k = \frac{12}{5}$$

Then eq. (i) becomes

$$p = \frac{12}{5} \frac{qr^2}{st^2}$$

Now for $q = -2$, $r = 4$, $s = 3$ and $t = -1$, we have

$$p = \frac{12}{5} \frac{(-2)(4)^2}{(3)(-1)^2} = -\frac{128}{5}$$

EXERCISE 3.5

1. If s varies directly as u^2 and inversely as v and $s = 7$ when $u = 3$, $v = 2$. Find the value of s when $u = 6$ and $v = 10$.
2. If w varies jointly as x , y^2 and z and $w = 5$ when $x = 2$, $y = 3$, $z = 10$. Find w when $x = 4$, $y = 7$ and $z = 3$.
3. If y varies directly as x^3 and inversely as z^2 and t , and $y = 16$ when $x = 4$, $z = 2$, $t = 3$. Find the value of y when $x = 2$, $z = 3$ and $t = 4$.
4. If u varies directly as x^2 and inversely as the product yz^3 , and $u = 2$ when $x = 8$, $y = 7$, $z = 2$. Find the value of u when $x = 6$, $y = 3$, $z = 2$.
5. If v varies directly as the product xy^3 and inversely as z^2 and $v = 27$ when $x = 7$, $y = 6$, $z = 7$. Find the value of v when $x = 6$, $y = 2$, $z = 3$.
6. If w varies inversely as the cube of u , and $w = 5$ when $u = 3$. Find w when $u = 6$.

3.4. K-Method

3.4(i) Use k -method to prove conditional equalities involving proportions.

If $a : b :: c : d$ is a proportion, then putting each ratio equal to k

$$\text{i.e., } \frac{a}{b} = \frac{c}{d} = k$$

$$\frac{a}{b} = k \text{ and } \frac{c}{d} = k$$

$$a = bk \text{ and } c = dk$$

Using the above equations, we can solve certain problems relating to proportions more easily. This method is known as k -method. We illustrate the k -method through the following examples.

Example 1: If $a : b = c : d$, then show that

$$\frac{3a + 2b}{3a - 2b} = \frac{3c + 2d}{3c - 2d}$$

Solution: $a : b = c : d$

$$\text{Let } \frac{a}{b} = \frac{c}{d} = k$$

$$\text{Then } a = bk \text{ and } c = dk$$

$$\text{To prove } \frac{3a + 2b}{3a - 2b} = \frac{3c + 2d}{3c - 2d}$$

$$\begin{aligned} \text{Now L.H.S} &= \frac{3a+2b}{3a-2b} = \frac{3kb+2b}{3kb-2b} = \frac{b(3k+2)}{b(3k-2)} \\ &= \frac{3k+2}{3k-2} \quad (\text{i}) \end{aligned}$$

$$\begin{aligned} \text{Also R.H.S} &= \frac{3c+2d}{3c-2d} = \frac{3kd+2d}{3kd-2d} = \frac{d(3k+2)}{d(3k-2)} \\ &= \frac{3k+2}{3k-2} \quad (\text{ii}) \end{aligned}$$

\therefore L.H.S = R.H.S

$$\text{i.e., } \frac{3a+2b}{3a-2b} = \frac{3c+2d}{3c-2d}$$

Example 2: If $a : b = c : d$, then show that

$$pa + qb : ma - nb = pc + qd : mc - nd$$

Solution: Let $\frac{a}{b} = \frac{c}{d} = k$, then $a = bk$ and $c = dk$

$$\begin{aligned} \text{L.H.S} = pa + qb : ma - nb &= \frac{pa + qb}{ma - nb} = \frac{pkb + qb}{mkb - nb} \\ &= \frac{b(pk + q)}{b(mk - n)} = \frac{pk + q}{mk - n} \end{aligned}$$

$$\begin{aligned} \text{R.H.S} = pc + qd : mc - nd &= \frac{pc + qd}{mc - nd} = \frac{pkd + qd}{mkd - nd} \quad (c = kd) \\ &= \frac{d(pk + q)}{d(mk - n)} = \frac{pk + q}{mk - n} \end{aligned}$$

$$\text{i.e., } pa + qb : ma - nb = pc + qd : mc - nd$$

Example 3: If $\frac{a}{b} = \frac{c}{d} = \frac{e}{f}$, then show that $\frac{a^3 + c^3 + e^3}{b^3 + d^3 + f^3} = \frac{ace}{bdf}$

Solution: Let $\frac{a}{b} = \frac{c}{d} = \frac{e}{f} = k$

$$\text{Then } \frac{a}{b} = k, \frac{c}{d} = k \text{ and } \frac{e}{f} = k$$

$$\text{i.e., } a = bk, c = dk \text{ and } e = fk$$

$$\text{To prove } \frac{a^3 + c^3 + e^3}{b^3 + d^3 + f^3} = \frac{ace}{bdf}$$

$$\begin{aligned} \text{Now L.H.S} &= \frac{a^3 + c^3 + e^3}{b^3 + d^3 + f^3} = \frac{(bk)^3 + (dk)^3 + (fk)^3}{b^3 + d^3 + f^3} \\ &= \frac{b^3k^3 + d^3k^3 + f^3k^3}{b^3 + d^3 + f^3} = k^3 \left(\frac{b^3 + d^3 + f^3}{b^3 + d^3 + f^3} \right) = k^3 \end{aligned}$$

$$\text{Also R.H.S.} = \frac{ace}{bdf} = \frac{(bk)(dk)(fk)}{bdf} = k^3 \frac{bdf}{bdf} = k^3$$

$$\therefore \text{L.H.S.} = \text{R.H.S.}$$

$$\text{i.e., } \frac{a^3 + c^3 + e^3}{b^3 + d^3 + f^3} = \frac{ace}{bdf}$$

Example 4: If $\frac{a}{b} = \frac{c}{d} = \frac{e}{f}$, then show that $\frac{a^2b + c^2d + e^2f}{ab^2 + cd^2 + ef^2} = \frac{a + c + e}{b + d + f}$

Solution: Let $\frac{a}{b} = \frac{c}{d} = \frac{e}{f} = k$

$$a = bk, c = dk, e = fk$$

$$\text{To prove } \frac{a^2b + c^2d + e^2f}{ab^2 + cd^2 + ef^2} = \frac{a + c + e}{b + d + f}$$

$$\text{L.H.S.} = \frac{a^2b + c^2d + e^2f}{ab^2 + cd^2 + ef^2}$$

$$= \frac{(bk)^2b + (dk)^2d + (fk)^2f}{(bk)b^2 + (dk)d^2 + (fk)f^2} = \frac{k^2b^3 + k^2d^3 + k^2f^3}{kb^3 + kd^3 + kf^3}$$

$$= \frac{k^2(b^3 + d^3 + f^3)}{k(b^3 + d^3 + f^3)} = k$$

$$\text{R.H.S.} = \frac{a + c + e}{b + d + f} = \frac{bk + dk + fk}{b + d + f}$$

$$= \frac{k(b + d + f)}{b + d + f} = k$$

$$\text{L.H.S.} = \text{R.H.S.}$$

$$\text{Thus, } \frac{a^2b + c^2d + e^2f}{ab^2 + cd^2 + ef^2} = \frac{a + c + e}{b + d + f}$$

EXERCISE 3.6

1. If $a : b = c : d$, ($a, b, c, d \neq 0$), then show that

$$(i) \quad \frac{4a - 9b}{4a + 9b} = \frac{4c - 9d}{4c + 9d} \quad (ii) \quad \frac{6a - 5b}{6a + 5b} = \frac{6c - 5d}{6c + 5d}$$

$$(iii) \quad \frac{a}{b} = \sqrt{\frac{a^2 + c^2}{b^2 + d^2}} \quad (iv) \quad a^6 + c^6 : b^6 + d^6 = a^3c^3 : b^3d^3$$

$$(v) \quad p(a + b) + qb : p(c + d) + qd = a : c$$

$$(vi) \quad a^2 + b^2 : \frac{a^3}{a + b} = c^2 + d^2 : \frac{c^3}{c + d}$$

$$(vii) \quad \frac{a}{a - b} : \frac{a + b}{b} = \frac{c}{c - d} : \frac{c + d}{d}$$

2. If $\frac{a}{b} = \frac{c}{d} = \frac{e}{f}$ ($a, b, c, d, e, f \neq 0$), then show that

$$(i) \quad \frac{a}{b} = \sqrt{\frac{a^2 + c^2 + e^2}{b^2 + d^2 + f^2}} \qquad (ii) \quad \frac{ac + ce + ea}{bd + df + fb} = \left[\frac{ace}{bdf} \right]^{2/3}$$

$$(iii) \quad \frac{ac}{bd} + \frac{ce}{df} + \frac{ea}{fb} = \frac{a^2}{b^2} + \frac{c^2}{d^2} + \frac{e^2}{f^2}$$

3.4(ii) Real life problems based on variation

Example 1: The strength “ s ” of a rectangular beam varies directly as the breadth b and the square of the depth d . If a beam 9cm wide and 12cm deep will support 1200 lb. What weight a beam of 12cm wide and 9cm deep will support?

Solution: By the joint variation, we have $s \propto bd^2$

$$i.e., \quad s = kbd^2 \qquad (i)$$

Put $s = 1200$, $b = 9$ and $d = 12$

$$k(9)(12)^2 = 1200$$

$$k = \frac{1200}{9 \times 144} = \frac{25}{27}$$

Put in eq. (i) $s = \frac{25}{27} bd^2$

Now for $b = 12$ and $d = 9$

$$s = \frac{25}{27} (12)(9)^2 = \frac{25(12)(9)(9)}{27} = 900 \text{ lb}$$

Example 2: The current in a wire varies directly as the electromotive force E and inversely as the resistance R . If $I = 32$ amperes, when $E = 128$ volts and $R = 8$ ohms. Find I , when $E = 150$ volts and $R = 18$ ohms.

Solution: In joint variation, we have $I \propto \frac{E}{R}$, i.e., $I = \frac{kE}{R}$ (i)

For $I = 32$, $E = 128$ and $R = 8$,

$$32 = \frac{k(128)}{8} \Rightarrow \frac{32 \times 8}{128} = k \Rightarrow k = 2$$

Put in eq. (i) $I = \frac{2E}{R}$.

Now for $E = 150$ and $R = 18$

$$I = \frac{2(150)}{18} = \frac{50}{3} \text{ amp.}$$

EXERCISE 3.7

- The surface area A of a cube varies directly as the square of the length l of an edge and $A = 27$ square units when $l = 3$ units.
Find (i) A when $l = 4$ units (ii) l when $A = 12$ sq. units.
- The surface area S of the sphere varies directly as the square of radius r , and $S = 16\pi$ when $r = 2$. Find r when $S = 36\pi$.
- In Hook's law the force F applied to stretch a spring varies directly as the amount of elongation S and $F = 32lb$ when $S = 1.6$ in. Find (i) S when $F = 50$ lb (ii) F when $S = 0.8$ in.
- The intensity I of light from a given source varies inversely as the square of the distance d from it. If the intensity is 20 candlepower at a distance of 12ft. from the source, find the intensity at a point 8ft. from the source.
- The pressure P in a body of fluid varies directly as the depth d . If the pressure exerted on the bottom of a tank by a column of fluid 5ft. high is 2.25 lb/sq. in, how deep must the fluid be to exert a pressure of 9 lb/sq. in?
- Labour costs c varies jointly as the number of workers n and the average number of days d . If the cost of 800 workers for 13 days is Rs. 286000, then find the labour cost of 600 workers for 18 days.
- The supporting load c of a pillar varies as the fourth power of its diameter d and inversely as the square of its length l . A pillar of diameter 6 inch and of height 30 feet will support a load of 63 tons. How high a 4 inch pillar must be to support a load of 28 tons?
- The time T required for an elevator to lift a weight varies jointly as the weight w and the lifting depth d varies inversely as the power p of the motor. If 25 sec. are required for a 4-hp motor to lift 500 lb through 40 ft, what power is required to lift 800 lb, through 120 ft in 40 sec.?
- The kinetic energy (K.E.) of a body varies jointly as the mass " m " of the body and the square of its velocity " v ". If the kinetic energy is 4320 ft/lb when the mass is 45 lb and the velocity is 24 ft/sec, determine the kinetic energy of a 3000 lb automobile travelling 44 ft/sec.

MISCELLANEOUS EXERCISE - 3

1. Multiple Choice Questions

Four possible answers are given for the following questions. Tick (✓) the correct answer.

- (i) In a ratio $a : b$, a is called
- | | |
|----------------|-------------------|
| (a) relation | (b) antecedent |
| (c) consequent | (d) None of these |

- (ii) In a ratio $x : y$, y is called
- (a) relation (b) antecedent
(c) consequent (d) None of these
- (iii) In a proportion $a : b :: c : d$, a and d are called,
- (a) means (b) extremes
(c) third proportional (d) None of these
- (iv) In a proportion $a : b :: c : d$, b and c are called
- (a) means (b) extremes
(c) fourth proportional (d) None of these
- (v) In continued proportion $a : b = b : c$, $ac = b^2$, b is said to be _____ proportional between a and c .
- (a) third (b) fourth
(c) means (d) None of these
- (vi) In continued proportion $a : b = b : c$, c is said to be _____ proportional to a and b .
- (a) third (b) fourth
(c) means (d) None of these
- (vii) Find x in proportion $4 : x :: 5 : 15$
- (a) $\frac{75}{4}$ (b) $\frac{4}{3}$
(c) $\frac{3}{4}$ (d) 12
- (viii) If $u \propto v^2$, then
- (a) $u = v^2$ (b) $u = kv^2$
(c) $uv^2 = k$ (d) $uv^2 = 1$
- (ix) If $y^2 \propto \frac{1}{x^3}$, then
- (a) $y^2 = \frac{k}{x^3}$ (b) $y^2 = \frac{1}{x^3}$
(c) $y^2 = x^2$ (d) $y^2 = kx^3$
- (x) If $\frac{u}{v} = \frac{v}{w} = k$, then
- (a) $u = wk^2$ (b) $u = vk^2$
(c) $u = w^2k$ (d) $u = v^2k$
- (xi) The third proportional of x^2 and y^2 is
- (a) $\frac{y^2}{x^2}$ (b) x^2y^2
(c) $\frac{y^4}{x^2}$ (d) $\frac{y^2}{x^4}$

(xii) The fourth proportional w of $x : y :: v : w$ is

(a) $\frac{xy}{v}$

(b) $\frac{vy}{x}$

(c) xyv

(d) $\frac{x}{vy}$

(xiii) If $a : b = x : y$, then alternando property is

(a) $\frac{a}{x} = \frac{b}{y}$

(b) $\frac{a}{b} = \frac{x}{y}$

(c) $\frac{a+b}{b} = \frac{x+y}{y}$

(d) $\frac{a-b}{x} = \frac{x-y}{y}$

(xiv) If $a : b = x : y$, then invertendo property is

(a) $\frac{a}{x} = \frac{b}{y}$

(b) $\frac{a}{a-b} = \frac{x}{x-y}$

(c) $\frac{a+b}{b} = \frac{x+y}{y}$

(d) $\frac{b}{a} = \frac{y}{x}$

(xv) If $\frac{a}{b} = \frac{c}{d}$, then componendo property is

(a) $\frac{a}{a+b} = \frac{c}{c+d}$

(b) $\frac{a}{a-b} = \frac{c}{c-d}$

(c) $\frac{ad}{bc}$

(d) $\frac{a-b}{b} = \frac{c-d}{d}$

2. Write short answers of the following questions.

(i) Define ratio and give one example.

(ii) Define proportion.

(iii) Define direct variation.

(iv) Define inverse variation.

(v) State theorem of componendo-dividendo.

(vi) Find x , if $6 : x :: 3 : 5$.

(vii) If x and y^2 varies directly, and $x = 27$ when $y = 4$. Find the value of y when $x = 3$.

(viii) If u and v varies inversely, and $u = 8$, when $v = 3$. Find v when $u = 12$.

(ix) Find the fourth proportional to 8, 7, 6.

(x) Find a mean proportional to 16 and 49.

(xi) Find a third proportional to 28 and 4.

(xii) If $y \propto \frac{x^2}{z}$ and $y = 28$ when $x = 7$, $z = 2$, then find y .

(xiii) If $z \propto xy$ and $z = 36$ when $x = 2$, $y = 3$, then find z .

(xiv) If $w \propto \frac{1}{v^2}$ and $w = 2$ when $v = 3$, then find w .

3. Fill in the blanks

- (i) The simplest form of the ratio $\frac{(x+y)(x^2+xy+y^2)}{x^3-y^3}$ is _____.
- (ii) In a ratio $x : y$; x is called _____.
- (iii) In a ratio $a : b$; b is called _____.
- (iv) In a proportion $a : b :: x : y$; a and y are called _____.
- (v) In a proportion $p : q :: m : n$; q and m are called _____.
- (vi) In proportion $7 : 4 :: p : 8$, $p =$ _____.
- (vii) If $6 : m :: 9 : 12$, then $m =$ _____.
- (viii) If x and y varies directly, then $x =$ _____.
- (ix) If v varies directly as u^3 , then $u^3 =$ _____.
- (x) If w varies inversely as p^2 , then $k =$ _____.
- (xi) A third proportional of 12 and 4, is _____.
- (xii) The fourth proportional of 15, 6, 5 is _____.
- (xiii) The mean proportional of $4m^2n^4$ and p^6 is _____.
- (xiv) The continued proportion of 4, m and 9 is _____.

SUMMARY

- A relation between two quantities of the same kind is called **ratio**.
- A **proportion** is a statement, which is expressed as equivalence of two ratios.

If two ratios $a : b$ and $c : d$ are equal, then we can write $a : b = c : d$

- If two quantities are related in such a way that increase (decrease) in one quantity causes increase (decrease) in the other quantity is called **direct variation**.
- If two quantities are related in such a way that when one quantity increases, the other decreases is called **inverse variation**.

- Theorem on proportions:

(1) **Theorem of Invertendo**

If $a : b = c : d$, then $b : a = d : c$

(2) **Theorem of Alternando**

If $a : b = c : d$, then $a : c = b : d$

(3) **Theorem of Componendo**

If $a : b = c : d$, then

(i) $a + b : b = c + d : d$

and (ii) $a : a + b = c : c + d$

(4) **Theorem of Dividendo**

If $a : b = c : d$, then

(i) $a - b : b = c - d : d$

(ii) $a : a - b = c : c - d$

(5) **Theorem of Componendo-dividendo**

If $a : b = c : d$, then

$$a + b : a - b = c + d : c - d$$

➤ A combination of direct and inverse variations of one or more than one variable forms **joint variation**.

➤ **K-Method,**

(a) If $\frac{a}{b} = \frac{c}{d}$,

then $\frac{a}{b} = \frac{c}{d} = k$ or $a = bk$ and $c = dk$

(b) If $\frac{a}{b} = \frac{c}{d} = \frac{e}{f} = k$, then $a = bk$, $c = dk$ and $e = fk$